

Selection of promising yeasts for starter culture development promoter of cocoa fermentation

Selección de levaduras promisorias para el desarrollo de un cultivo iniciador promotor de la fermentación del cacao

Jenifer Criollo-Nuñez¹⁻⁴ Angélica P Sandoval- Aldana² German Bolívar³ Cristina Ramírez Toro⁴

¹ Corporación Colombiana de Investigación Agropecuaria (AGROSAVIA), Centro de Investigación Nataima, Ibagué Tolima, Colombia.

² Universidad del Tolima, Facultad de Ingeniería Agronómica, Ibagué Tolima, Colombia.

³ Universidad del Valle, Facultad de Ciencias Naturales y Exactas, Cali Valle del Cauca, Colombia.

⁴ Universidad del Valle, Facultad de Ingeniería, Cali Valle del cauca, Colombia.

Abstract

The selection of promising microorganisms that meet the requirements of a fermentation process is the most important stage in the development of microbiological starters. With the objective of selecting potential strains to be integrated as microbiological starters in cocoa fermentations, which promote biochemical dynamics, improving their sensory quality, the resistance of six yeasts (GRAS), isolated from cocoa fermenting masses in the departments, was evaluated. from Tolima, Huila and Antioquia, two commercial yeasts ATCC 28253 and ATCC 36633 selected for their aromatic production, subjecting them to resistance tests, at different concentrations of soluble solids (10, 15 and 20 °Brix), ethanol (5, 10%), acetic acid (pH 3, 4, 5, 6, and 7) and temperature (30, 40 and 50 °C), which simulated the biochemical dynamics of cocoa fermentation, quantifying its resistance expressed in viable cells. Aromatic production was measured by sensory evaluation. The yeasts *Wickerhamomyces anomalus* and *Pichia kudriavzevii* were selected for their ability to transform substrate into products of interest during the fermentation of cocoa pulp, production of specific aromas and resistance to adverse biochemical conditions typical of cocoa fermentation.

Resumen

La selección de microorganismo promisorios que atiendan los requerimientos de un proceso fermentativo es la etapa más importante en el desarrollo de iniciadores microbiológicos. Con el objetivo de seleccionar las cepas potenciales para ser integradas como iniciadores microbiológicos en fermentaciones de cacao, que promuevan la dinámica bioquímica, mejorando su calidad sensorial, se evaluó la resistencia de seis levaduras (GRAS), aisladas de masas fermentantes de cacao en los departamento del Tolima, Huila y Antioquia, dos levaduras comerciales ATCC 28253 y ATCC 36633 seleccionadas por su producción aromática, sometiénolas a pruebas de resistencia, a diferentes concentraciones de sólidos solubles (10, 15 y 20 °Brix), etanol (5, 10 %), ácido acético (pH 3, 4, 5, 6, y 7) y temperatura (30, 40 y 50 oC), que simularon la dinámica bioquímica de la fermentación del cacao, cuantificando su resistencia expresada en células viables. La producción aromática fue medida mediante evaluación sensorial. Se seleccionaron las levaduras *Wickerhamomyces anomalus* y *Pichia kudriavzevii* por su capacidad de transformar sustrato en productos de interés durante la fermentación de pulpa de cacao, producción de aromas específicos y resistencia a condiciones bioquímicas adversas propias de la fermentación del cacao.

Keywords: yeasts, cocoa fermentation, microbiological Starter, Biochemical dynamics, Quality.

Palabras clave: levaduras, Fermentación cacao, Iniciador microbiológico, Dinámica bioquímica, Calidad.

How to cite?

Criollo-Nuñez, J., Sandoval-Aldana, A., Bolívar, G., Ramírez, C. Selection of promising yeasts for the development of a microbiological starter that promotes cocoa fermentation *Ingeniería y Competitividad*, 2024, 26(2)e-20413296.

<https://doi.org/10.25100/iyc.v26i2.13296>

Recibido: 19-10-23 Aceptado: 20-04-24

Correspondence:

jcriollo@agrosavia.co

This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike4.0 International License.



Conflict of interest: none declared



Why was it carried out?

The use of microbiological starters in the food industry to improve processes and the quality of the finished product demands the selection of resistant microorganisms with the potential to transform substrate into metabolites of interest. Seeking to promote the biochemical dynamics of cocoa fermentation, the potential of eight yeasts was evaluated against different biochemical conditions that simulated the cocoa fermentation process to develop a microbiological starter that will improve the biochemical dynamics of fermentation and the sensory quality of cocoa.

What were the most relevant results?

The yeasts *Wickerhamomyces anomalus* and *Pichia kudriavzevii* are promising strains for the development of microbiological starters due to their ability to transform substrate into products of interest during the fermentation of cocoa pulp, with the production of specific aromas and resistance to adverse biochemical conditions typical of the cocoa fermentation. The selected microorganisms have high feasibility of application as a starter culture promoter of fermentation, which can be used individually, by consortium, and in mixed culture.

What do these results provide?

The identification of yeast strains promising for the development of a microbiological starter that promotes the dynamics of cocoa fermentation that improves the sensory characteristics of chocolate.

Graphical Abstract



Introduction

The genetic diversity of cacao and differences in its biochemical characteristics generate a wide range of fermentation conditions. This is an important stage in the cacao postharvest, where a series of biochemical transformations induced by microbial action (microbial succession) develop, modifying the substrate in the pulp into substances of interest that modify the sensory profile of cocoa beans.

The cacao microbiota, which has been shown to intervene positively during fermentation, is composed of yeasts, lactic acid bacteria (LAB), and acetic acid bacteria (AAB) (1). When the fermentation stage is prolonged for more than four days, bacilli, enterobacteria, and filamentous fungi, desired or not, can also participate in the cocoa fermentation process (2). Yeasts have been related to the production of volatile organic compounds generators of typical aromas of some fermented foods, associated with the production of endogenous enzymes that induce the generation of chemical precursors of flavor such as free amino acids, peptides, and reducing sugars (3–5).

The application of starter cultures in fermentation has been studied to improve microbial dynamics and biochemical transformations in the process, reduce time, and improve the sensory quality of the bean. Recently, interest has grown in using yeasts as starter cultures for cocoa fermentation, especially species of the *Saccharomyces*, *Pichia*, *Kluyveromyces*, *Candida*, and *Torulaspota* genera (6,7). Furthermore, some studies evaluated the application of mixed starters composed of yeast and LAB (5,8). The great capacity to assimilate citric acid, reducing sugars, pectinolytic activity, resistance to stress conditions, production of volatile compounds, and antagonistic capacity against pathogens has directed research towards using these microorganisms as a biotechnological strategy to promote sensory quality in cocoa (9,10). This requires the selection of microorganisms resistant to the conditions of the fermentation process, guaranteeing a high transformation of substrates and generating products of interest that modify the sensorial quality of cocoa to show a greater aroma or flavor. For this reason, there is interest in isolating promising microorganisms to be inoculated during the cocoa fermentation phase. This research aims to select yeasts according to their morphological characterization, behavior during the fermentation process, and the production of aromas through a sensory evaluation, with potential application for optimizing the fermentation phase and sensory quality of cocoa.

Methodology

Microorganisms

Eight yeasts were evaluated, six isolated from fermentative masses of cocoa-producing farms in the departments of Huila, Tolima, and Antioquia, Colombia, including *Hanseniaspora opuntiae* (Y01), *Saccharomyces cerevisiae* (Y02), *Pichia kudriavzevii* (Y03), *Debaryomyces hansenii* (Y04), *Aureobasidium pullulans* (Y07), and *Wickerhamomyces anomalus* (Y08) of the microorganisms working collection of Agrosavia with interest in nutrition and animal health. The other two, ATCC 36633 *Candida versatilis* (Etchells and Bell) Meyer and Yarrow (Y05) and ATCC 28253 *Zygosaccharomyces rouxii* (Boutroux) Milenrama (Y06), were certified commercial yeast strains selected for their production of aromatic compounds.

Yeast tolerance tests to simulated cocoa fermentation process conditions

The microorganisms were subjected to resistance tests, inoculating 1,000 μL of the microorganism suspension with turbidity equal to McFarland tube 3 in 9,000 μL of water peptone at 0.1 % in 20 mL test tubes. Prior to inoculation, the malt extract broths were conditioned at soluble solids concentrations of 10, 15, and 20 °Brix, ethyl alcohol concentrations at 5 and 10 %, acetic acid with pH of 3, 4, 5, 6, and 7, and temperatures of 30, 40, and 50 °C in the broths. The cultures were incubated for 48 h under constant agitation at 120 rpm, quantifying viable cells in decimal dilutions 10^{-4} , 10^{-5} , and 10^{-6} at the

end of the incubation, according to the methodology proposed by Jurado Gámez et al. (11) and Lozano Tovar et al. (12). The uninoculated culture medium was used as a control.

Substrate utilization test

The yeasts were inoculated individually in 300 mL of sterile cocoa pulp medium with a 3:1 pulp:water ratio, adding 15 mL of suspension of the microorganism with turbidity equal to McFarland tube 3 and incubating under constant agitation at 1,200 rpm and temperature of 30 °C for 35 h, using the adapted methodology of Briceño Vélez (13). The fermentations were carried out in triplicate, taking aliquots at 0, 5, 10, 15, 20, 25, 30, and 35 h of fermentation, quantifying the content of soluble solids, reducing sugars, pH, titratable acidity, percentage of ethyl alcohol and viable cells, following the methodologies detailed as follows.

Soluble solids, ethyl alcohol, pH, and titratable acidity: The aliquots at different fermentation times were analyzed for total soluble solids in °Brix, taking 1 mL of the sample and measuring in a portable refractometer PAL-1 (3810) (Atago Co. Ltd, Tokyo, Japan). The percentage of ethyl alcohol was registered with a digital refractometer PAL-34S (4434) (Atago Co. Ltd, Tokyo, Japan) in percentage (%). The volatile acidity or pH in the sample was quantified in an aliquot of 10 mL using a potentiometer (Thermo Scientific). Total acidity was estimated in percentage (%) by reducing the acidity present in the sample with a basic 0.1 N NaOH solution (14).

Reducing sugars: Dilutions of 1:1000 v/v were made to quantify fermentable sugars in the cocoa pulp medium. All samples were homogenized for 5 min and passed through a 0.22 µm millipore filter. The content of reducing sugars was calculated using the 3,5 dinitro salicylic acid (DNS) method, a compound that, in the presence of heat, is reduced by the reducing sugars present, developing a color change. It was measured in g/L using a spectrophotometer SQ2800 Unico at a wavelength of 575 nm, estimating the reducing units present in sugars based on a glucose standard curve (15).

Viable cells: The viability or viable microorganisms count was estimated by diluting 1 mL of the aliquot at different fermentation times in 9 mL of 0.01% peptone water. Decimal dilutions were made up to 10⁻⁶; the last three dilutions were cultured in malt extract medium placed in Petri dishes and in superficial extension with glass beads. The dishes were incubated at 35°C for 48 h, counting the colonies on the plate, which were multiplied by the dilution factor and by 10, expressing the results in colony-forming units per milliliter (CFU/mL) (16).

Sensory analysis to detect aroma production in cocoa pulp

In 150 mL of cocoa pulp medium in a 3:1 pulp:water ratio, 15 mL of microorganism suspension was inoculated at turbidity equal to McFarland tube 3, incubating at low oxygen pressures for 48 h, in triplicate. The 20 mL aliquots were evaluated by trained tasters using the descriptive olfactory profile method. Samples were analyzed in terms of odor descriptors (sour, sweet, floral, fruity, alcohol, herbal, and yeasty), using an ordinal scale from 0 to 10, where 0 means absence, 1 to 2 low intensity, 3 to 5 medium intensity, 6 to 8 high intensity, and 9 to 10 very high intensity. The panel was also asked about their three preferred samples concerning aroma formation (17).

Statistical analysis

The data on resistance to yeast fermentation dynamics and substrate utilization were normalized. Subsequently, parametric analysis of variance (ANOVA) tests were performed, followed by the multiple comparison test for differences between the Tukey means with a 95% confidence level using the RStudio statistical software.

Results and discussion

Tolerance to different pH conditions

Cocoa fermentation shows a microbiological dynamic that generates biochemical changes in parameters such as pH. Figure 1 shows the tolerance expressed in the viability of the yeasts studied when subjected to different pH conditions, taking as reference what was expressed by authors such as Bilenler et al. (18), who reported as the minimum required viability for microbiological starters applied in foods a value of 10^6 CFU/g.

The yeast *Hanseniaspora opuntiae* Y01 reported the highest viability at a pH of 3, which was drastically reduced when subjected to other pH values of the biochemical dynamics, reaching values below 6 log at pH values of 5 and 7. This behavior is analogous to the cocoa microbial diversity study results, where this species was isolated only at the beginning of fermentations (10), and its detection was not possible at later fermentation times.

The viability of *S. cerevisiae* Y02, *Z. rouxii* Y06, and *A. pullulans* Y07 was only affected when subjected to pH values of 7, 6, and 3, respectively, without registering statistically significant differences in the other pH values studied. Yeast *W. anomalus* Y08 did not report statistically significant differences in pH, maintaining its viability among the highest microbiological density levels compared to the other yeasts in the study. In this way, its growth capacity at different pH conditions and adaptability to different biochemical conditions were demonstrated, positioning it as a potential yeast to be incorporated into biochemically dynamic fermentative processes without its viability being affected (19).




Figure 1. Yeast tolerance test to different pH growing conditions.

Tolerance to different temperature conditions

Figure 2 shows the tolerance of the yeasts under study to different temperature conditions of cocoa fermentation dynamics. Yeasts must be resistant to temperatures above $45\text{ }^{\circ}\text{C}$ to guarantee the death of the seed embryo, so this factor is decisive in the selection of the microorganism (20,21).

The high-temperature tolerance of the yeast *P. kudriavzevii* (Y03) was highlighted, being the only viable strain exposed to temperature conditions of 50 °C, maintaining its viability above 6 Log, coinciding with the report of its effectiveness in growth capacity and high-temperature ethyl alcohol production from sugar cane juice, corn stalk hydrolyzate and cassava hydrolyzate (22). Additionally, this species recorded the highest viability at all the temperatures studied. Strains *D. hansenii* (Y04), *Z. rouxii* (Y06), *A. pullulans* (Y07), and *W. anomalus* (Y08) recorded statistically significant differences at a confidence level $P > 0.05$ for the temperature factor being its viability affected by the thermal level (23).




Figure 2. Yeast tolerance test to different temperature (°C) growing conditions.

Tolerance to different soluble solids conditions

The genetic variability of cacao materials and their bromatological composition demand a starter culture for cocoa fermentation, with the capacity to tolerate differences in the soluble solids content of its pulp without affecting its viability. The results of the tolerance test of the studied yeasts are recorded in Figure 3. Strain *W. anomalus* (Y08) registered the highest viability at 10 and 15 °Brix, with 8 log and 7 log in culture media with soluble solids contents of 20 °Brix (Figure 3). Its osmotolerance was highlighted, also recorded by the strain *Candida versatilis* (Y05), which has been isolated from apiaries and used in soybean fermentation processes, emphasizing its capacity to convert sugars into ethyl alcohol, organic acids metabolism, and volatile compounds formation, describing it as a tolerant and adaptive yeast.




Figure 3. Yeast tolerance test to different soluble solids (°Brix) growing conditions. Tolerance to different ethyl alcohol conditions

Figure 4 shows the tolerance of the yeasts under study to 5 and 10 % ethyl alcohol concentrations.




Figure 4. Yeast tolerance test to different ethyl alcohol (%) growing conditions.

No statistically significant differences were recorded in the viability of the yeasts when they were exposed to 5% ethyl alcohol. However, with the increase in ethyl alcohol concentration, yeast strains such as Y01, Y04, and Y07 were strongly affected in their viability. The resistance of strains Y05, Y03, Y02, and Y06 to high concentrations of ethyl alcohol in the culture was highlighted, registering viabilities above 7 log, which, in strains such as *P. kudriavzevii* (Y03), demonstrated its ability to withstand adverse conditions during cocoa fermentation by remaining viable under conditions of high ethyl alcohol concentration and temperature. Therefore, it can be considered a promising microorganism to be used as a starter culture in cocoa fermentation.

The tolerance tests allowed generating a first screening of the yeasts studied to select a starter culture to improve the microbial dynamics during fermentation and the sensory quality of cocoa beans. The yeast strain *H. opuntiae* (Y01) was discarded as it registered low tolerance in factors such as pH, soluble solids content, temperature, and ethyl alcohol concentration of 10%, reflected in its low viability and resistance to the biochemical dynamics of cocoa fermentation.

Substrate utilization consumption test

The capacity and speed of transforming macromolecules into a substance of interest are other selection parameters that the starter culture must meet. Yeasts interact in the first phase of cocoa bean fermentation under acidic conditions and low oxygen pressures, generating various transformations, as stated by Delgado-Ospina et al. (24). Ethyl alcohol is produced from citric acid and carbohydrates (sucrose, glucose, and fructose), present in the mucilage of ripe cacao fruits, which partially diffuses into the cotyledons of the cacao bean, oxidizing later on due to the effect of acetic acid bacteria (3,25). Figure 5 shows the viability, substrate utilization capacity, and acid and alcohol production of the yeasts studied. Yeast strains Y02 and Y03 showed the culmination of their exponential phase at 10 h of fermentation in cocoa pulp medium, which for strains Y07 and Y08 was reported at 25 h of fermentation. Yeast *Zygosaccharomyces rouxii* (Y06) recorded a prolonged stationary phase by maintaining its highest viability between 15 and 25 h of fermentation. These results show the specificity in the viable behavior of yeasts of different genera on the same substrate (23).

The content of soluble solids showed a decreasing behavior for the five preselected yeasts in the 35 h of fermentation, with the highest decrease found in the musts of *Z. rouxii* (Y06) and *S. cerevisiae* (Y02). A similar evolutionary profile was recorded in the content of reducing sugars generated by the utilization of carbohydrates and glucose, which were hydrolyzed from the sucrose of cacao pulp by invertase activity. This explains the increase reported in the reducing sugar content of *W. anomalus* (Y08), *P. kudriavzevii* (Y03), and *S. cerevisiae* (Y02). During the first five hours of fermentation, they showed a greater availability of fermentable sugars or monosaccharides that are subsequently transformed into ethyl alcohol as a primary product and carbon dioxide and glycerol as secondary products (10,25).




Figure 5. Substrate utilization dynamics by preselected yeasts. Viability (A), Ethylic alcohol production (B), Soluble solids dynamics (C), Reducing sugars (D), Volatile acidity (E), and Titratable acidity (F).

The pH presented a decreasing behavior during the first 25 h of fermentation with the five preselected yeasts, which demonstrates the heterofermentative capacity of the yeasts under study, capable of producing organic acids such as acetic acid and succinic acid (26,27), compounds probably responsible for the increase in the total acidity of the medium. After 25 h of fermentation, an increase in pH was recorded, possibly generated by reducing the main carbon source (carbohydrates) and promoting the consumption of acids in the medium.

No statistically significant differences were recorded in the ethyl alcohol production of *P. kudriavzevii* Y03, *W. anomalus* Y08, *S. cerevisiae* Y02, and *Z. rouxii* Y06. However, *P. kudriavzevii* Y03 and *W. anomalus* Y08 were those with the highest production of the metabolite of interest, recording a lower consumption of reducing sugars and total acidity, evidencing a higher capacity to transform sugars and acids present in the cacao pulp into alcohol, required for the death of the embryo and subsequent diffusion in cacao beans during the aromatic development of the beans (6,20).

Production of yeast aromas in cacao pulp musts

The production of aromas during cacao fermentation has been mainly associated with the microbiological consortium of yeasts in the process (10,25,28,29). It is subjected to factors such as the nutrient concentration of the substrate, biochemical conditions of the process, and yeast genetics that, in turn, intervene in the production of enzymes and generation of metabolites of interest, as expressed by Kongor et al. (30) and Moreira et al. (31). The aroma production test in the cacao pulp substrate of the yeasts under study is shown in Figure 6 through the aroma profile of the musts.




Figure 6. Sensory profile of aroma production of the yeasts under study.

The qualification of fine and aromatic cocoa requires the presence of specific notes in its sensory profile, highlighting yeasts such as *P. kudriavzevii* Y03, *S. cerevisiae* Y02, *Z. rouxii* Y06, and *W. anomalus* Y08, which recorded the highest intensity in the fruit aroma of the musts studied with average values of 7, 6, 5 and 5, respectively. The floral aroma was

detected in greater intensity in the aromatic production of yeast *Z. rouxii* Y08, a species that also recorded herbal and honey notes in its profile that were not found in the fermented musts of the other yeasts studied, agreeing with the high production of ethyl acetate reported for this species and that has been related to fruity notes (32). Phenylethyl alcohol and 3-methyl-1-butanol acetate detected in rice wine fermentation are associated with fruity and floral notes; hence, this species stands out for its aromatic production (33). The presence of alcohol was detected at a higher intensity in the fermented musts of yeasts *P. kudriavzevii* Y03, *Z. rouxii* Y08, and *S. cerevisiae* Y02. Since this compound is required for the death of the embryo during cocoa fermentation, its greater presence can favor the release of compounds that act as precursors of organoleptic molecules. Yeasts like *D. hansenii* Y04 and *C. versatilis* Y05 recorded in their profile unpleasant aromas (off-flavors) in intensities of 3 and 2.5, respectively, which were described as overfermentation or decomposed fruit. These, together with the low production of specific aromas and alcohol and less resistance to biochemical dynamics compared to the other yeasts studied, allow their exclusion from the yeast selection process as a starter culture for cocoa fermentation.

Conclusions

Yeasts *W. anomalus* (Y08) and *P. kudriavzevii* (Y03) were selected for their ability to transform substrate into products of interest during the fermentation of cacao pulp, with the production of specific aromas and resistance to adverse biochemical conditions typical of cacao fermentation. The selected microorganisms have a high feasibility of application as a fermentation promoter starter culture that can be used individually, by consortium, and as a mixed culture.

Acknowledgements

The authors thank the Department of Tolima and Minciencias for funding this research through Call no. 755 entitled "Training of high-level human capital for the Department of Tolima," Agreement 2078 between the Government of Tolima and Universidad del Tolima with the title "Development of technological solutions for cocoa production and processing," Corporación colombiana de investigación agropecuaria (Agrosavia) within the framework of a Memorandum of Understanding with Universidad del Tolima S464 and Universidad del Valle through the project "Development of a biotechnological strategy to improve fermentation dynamics and quality" [code 528-2019]. This document is the result of the research project funded by the Government of Tolima and Universidad del Tolima [Agreement No. 2078] entitled "Development of technological solutions for cocoa production and processing" [Internal Code 130617].

References

1. Apriyanto M, Novitasari R. During Fermentation, Microbiology and Biochemistry of the Cocoa Bean. International Journal of Food Science and Agriculture [Internet]. [cited 2022 Apr 13];2021(4):688–91. Available from: <http://www.hillpublisher.com/journals/jsfa/>
2. Figueroa-Hernández C, Mota-Gutierrez J, Ferrocino I, Hernández-Estrada ZJ, González-Ríos O, Cocolin L, et al. The challenges and perspectives of the selection of starter cultures for fermented cocoa beans. Int J Food Microbiol [Internet]. 2019 [cited 2021 Apr 6];301:41–50. Available from: <https://doi.org/10.1016/j.ijfoodmicro.2019.05.002>
3. Loviso CL, Libkind D. Synthesis and regulation of flavor compounds derived from brewing yeast: Esters. Rev Argent Microbiol. 2018 Oct 1;50(4):436–46.

4. Venturini Copetti M. Yeasts and molds in fermented food production: an ancient bioprocess. *Curr Opin Food Sci.* 2019 Feb 1;25:57–61.
5. Crafac M, Mikkelsen MB, Saerens S, Knudsen M, Blennow A, Lowor S, et al. Influencing cocoa flavour using *Pichia kluyveri* and *Kluyveromyces marxianus* in a defined mixed starter culture for cocoa fermentation. *Int J Food Microbiol.* 2013;167(1):103–16.
6. Batista KA, Fernandes KF. Development and optimization of a new culture media using extruded bean as nitrogen source. *MethodsX* [Internet]. 2015;2:154–8. Available from: <http://dx.doi.org/10.1016/j.mex.2015.03.001>
7. Melo Pereira G, Soccol V, Soccol C. Current state of research on cocoa and coffee fermentations. *Curr Opin Food Sci* [Internet]. 2016 [cited 2023 May 8];7:50–7. Available from: <https://www.sciencedirect.com/science/article/pii/S2214799315001307>
8. Crafac M, Keul H, Eskildsen CE, Petersen MA, Saerens S, Blennow A, et al. Impact of starter cultures and fermentation techniques on the volatile aroma and sensory profile of chocolate. *Food Research International.* 2014;63(November):306–16.
9. Gutiérrez Ríos, H.G., Suárez Quiroz, M.L., Hernández Estrada, Z.J., Castellanos Onorio, O.P., Alonso Villegas, R., Rayas Duarte, P., Cano Sarmiento, C., Figueroa Hernández, C. Y., González-Rios, O. (2022). Yeasts as producers of flavor precursors during cocoa bean fermentation and their relevance as starter cultures: A review. *Fermentation.* 8, 331. <https://doi.org/10.3390/fermentation8070331>
10. Maicas S. The Role of Yeasts in Fermentation Processes. *Microorganisms* 2020, Vol 8, Page 1142 [Internet]. 2020 Jul 28 [cited 2022 Apr 24];8(8):1142. Available from: <https://www.mdpi.com/2076-2607/8/8/1142/htm>
11. Jurado Gámez H, Ramírez C, Aguirre D. Cinética de fermentación de *Lactobacillus plantarum* en un medio de cultivo enriquecido como potencial probiótico. *Veterinaria y Zootecnia.* 2013 Dec 20;Vol 7 No.2:37–53.
12. Lozano Tovar MD, Tibasosa G, González CM, Ballestas Alvarez K, Lopez Hernandez MDP, Rodríguez Villamizar F. Isolation and identification of microbial species found in cocoa fermentation as microbial starter culture candidates for cocoa bean fermentation in Colombia. *Pelita Perkebunan (a Coffee and Cocoa Research Journal).* 2020;36(3):236–48.
13. Briceño Vélez DA. Efecto de las bacterias lácticas microencapsuladas en la calidad microbiológica y en las propiedades fisicoquímicas de salchichas cocidas [Maestría en Ingeniería de alimentos]. [Santiago de Cali]: Universidad del Valle; 2020.
14. AOAC. Official Methods of Analysis [Internet]. Vol. 21st Edition. 2019 [cited 2022 Apr 21]. Available from: <https://www.aoac.org/official-methods-of-analysis-21st-edition-2019/>
15. Dumitru M, Tabuc C, Sorescu I, Vasilachi A, Hăbeanu M, Petre S, et al. Researches concerning the level of fermentable sugars from feed materials in relation with cellulase hydrolysis by carbohydrase enzyme. *Scientific Bulletin Series F Biotechnologies.* 2018;XXII.
16. Nuñez JC, Toro CR, Bolívar G, Aldana AS, Tovar MDL. Effect of microencapsulated inoculum of *Pichia kudriavzevii* on the fermentation and sensory quality of cacao



- CCN51 genotype. *J Sci Food Agric* [Internet]. 2023 Jan 6 [cited 2023 Jan 15]; Available from: <https://onlinelibrary.wiley.com/doi/full/10.1002/jsfa.12433>
17. Nutricional–IIN MDI de I. Guía para la evaluación sensorial de alimentos. academia.edu [Internet]. [cited 2024 Jan 20];2007. Available from: <https://www.academia.edu/download/61102947/Guia-para-la-evaluacion-sensorial-de-alimentos20191102-104797-1le28kv.pdf>
 18. Bilenler T, Karabulut I, LWT KC. Efectos de los cultivos iniciadores encapsulados sobre las propiedades microbianas y fisicoquímicas de salchichas (sucuks) producidas tradicionalmente y tratadas térmicamente. Elsevier [Internet]. 2016 [cited 2022 Jul 23]; Available from: <https://www.sciencedirect.com/science/article/pii/S0023643816305655>
 19. Naumov G, Naumova ES, Schnürer J, Naumov GI, Naumova ES. Caracterización genética de la levadura no convencional *Hansenula anomala*. *Investigación en Microbiología* [Internet]. 2001 [cited 2022 Jul 23];152:551–62. Available from: <https://www.sciencedirect.com/science/article/abs/pii/S0923250801012293>
 20. Calvo AM, Botina BL, García MC, Cardona WA, Montenegro AC, Criollo J. Dynamics of cocoa fermentation and its effect on quality. *Sci Rep* [Internet]. 2021 Dec 1 [cited 2022 Mar 23];11(1):16746. Available from: [/pmc/articles/PMC8373873/](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8373873/)
 21. De Brito ES, García NHP, Gallão MI, Cortelazzo AL, Fevereiro PS, Braga MR. Structural and chemical changes in cocoa (*Theobroma cacao* L) during fermentation, drying and roasting. *J Sci Food Agric*. 2001;81(2):281–8.
 22. Chamnipa N, Thanonkeo S, Klanrit P, Thanonkeo P. The potential of the newly isolated thermotolerant yeast *Pichia kudriavzevii* RZ8-1 for high-temperature ethyl alcohol production. *Braz J Microbiol* [Internet]. 2018 Apr 1 [cited 2022 Jul 21];49(2):378–91. Available from: <https://pubmed.ncbi.nlm.nih.gov/29154013/>
 23. Frazier WC, Mesthoff DC. *Microbiología de alimentos* [Internet]. En lengua española. Vol. 4. Zaragoza España: ACRIBIA; 1993 [cited 2022 Jul 20]. 1–681 p. Available from: <https://iselavictoria06wordpress.files.wordpress.com/2019/04/l33.pdf>
 24. Delgado-Ospina J, Acquaticci L, Molina-Hernandez JB, Rantsiou K, Martuscelli M, Franks Kamgang-Nzekoue A, et al. Exploring the capability of yeasts isolated from Colombian fermented cocoa beans to form and degrade biogenic amines in a lab-scale model system for cocoa. *mdpi.com* [Internet]. 2020 [cited 2022 Apr 17]; Available from: <https://www.mdpi.com/934894>
 25. Ruta LL, Farcasanu IC. Coffee and yeasts: From flavor to biotechnology. *Fermentation*. 2021;7(1).
 26. Romanens E, Freimüller Leischfeld S, Volland A, Stevens M, Krähenmann U, Isele D, et al. Screening of lactic acid bacteria and yeast strains to select adapted anti-fungal co-cultures for cocoa bean fermentation. *Int J Food Microbiol*. 2019;290(October):262–72.
 27. Schwan RF, Pereira GV de M, Fleet GH. Microbial activities during cocoa fermentation. *Cocoa and coffee fermentations*. 2014;(January 2014):129–92.
 28. De Vuyst L, Leroy F. Functional role of yeasts, lactic acid bacteria and acetic acid bacteria in cocoa fermentation processes. *FEMS Microbiol Rev*. 2020;(May):432–53.
 29. Fonseca Blanco JD, López Hernandez M del P, Ortiz Galeano LS, Criollo Nuñez J, Lozano Tovar MD. Effect of addition of a specific mixture of yeast, lactic and acetic bacteria in the fermentation process to improve the quality and flavor of cocoa



- beans in Colombia. *ccrjournal.com* [Internet]. 2020 [cited 2022 Apr 17];36(2):154–72. Available from: <https://ccrjournal.com/index.php/ccrj/article/view/438>
30. Kongor JE, Hinneh M, de Walle D Van, Afoakwa EO, Boeckx P, Dewettinck K. Factors influencing quality variation in cocoa (*Theobroma cacao*) bean flavour profile - A review. *Food Research International* [Internet]. 2016;82:44–52. Available from: <http://dx.doi.org/10.1016/j.foodres.2016.01.012>
 31. Moreira I, Costa J, Vilela L, Lima N, Santos C, Schwan RF. Influence of *S. cerevisiae* and *P. kluyveri* as starters on chocolate flavour. *J Sci Food Agric*. 2021;101:4409–19.
 32. Zha M, Sun B, Wu Y, Yin S, Wang C. Improving flavor metabolism of *Saccharomyces cerevisiae* by mixed culture with *Wickerhamomyces anomalus* for Chinese Baijiu making. *J Biosci Bioeng*. 2018 Aug 1;126(2):189–95.
 33. Wu Y, Chen X, Fang X, Ji L, Tian F, Yu H, et al. Isolation and Identification of Aroma-producing Yeast from Mackerel Fermentation Broth and Its Fermentation Characteristics. <https://doi-org.recursos.agrosavia.co/101080/1049885020211988016> [Internet]. 2021 [cited 2022 Sep 30];30(10):1264–80. Available from: <https://www-tandfonline-com.recursos.agrosavia.co/doi/abs/10.1080/10498850.2021.1988016>