

# State of the Art of Agricultural Water Footprint in Colombia

## Estado del arte de la huella hídrica agrícola en Colombia

Andrés Felipe Góngora-Duarte<sup>1</sup>   Juan Manuel Trujillo-González<sup>2</sup> 

<sup>1</sup>Ingeniero Agroindustrial, Maestría en Gestión Ambiental Sostenible, Facultad de Ciencias Básicas e Ingeniería, Universidad de los Llanos Universidad de los Llanos Campus Barcelona, Villavicencio Colombia.

<sup>2</sup>Instituto de Ciencias Ambientales de la Orinoquia Colombiana ICAOC, Universidad de los Llanos Campus Barcelona, Villavicencio Colombia.

### Abstract

**Introduction:** The water footprint is an environmental metric that quantifies the total volume of water used in the production of goods or services. In agriculture, it is measured in cubic meters per ton of product (m<sup>3</sup>/ton) and comprises three components: green (rainwater), blue (surface or groundwater), and grey (polluted water). This metric helps assess the water-use efficiency of crops.

**Objectives:** This study aimed to examine water use in Colombian agriculture through a systematic review, focusing on water resource management and its relevance for guiding decisions on the optimal geographic distribution of crops based on their water requirements.

**Materials and Methods:** A systematic review was conducted, analyzing 37 scientific studies on agricultural water footprints in Colombia, covering a total of 44 crops across various production regions. Reported values for the water footprint per crop were examined, along with the proportional distribution of green, blue, and grey water components.

**Results:** The most frequently studied crops were vegetables (20%), coffee and cocoa (19%), fruits (17%), flowers (12%), cereals (10%), and others (22%). On average, 72% of the total water use corresponded to green water, 15% to blue water, and 13% to grey water, indicating a high reliance on rainwater in Colombian agriculture.

**Conclusions:** The findings underscore the importance of evidence-based water planning to enhance water-use efficiency in Colombia's agricultural sector, thereby supporting sustainable production and improved resource management.

**Keywords:** Water Footprint, Agriculture, virtual water, sustainability, water resources.

### Resumen

**Introducción:** La huella hídrica es una métrica ambiental que cuantifica el volumen total de agua utilizado en la producción de bienes o servicios. En la agricultura, esta se mide en m<sup>3</sup> por tonelada de producto (m<sup>3</sup>/ton) y se divide en tres componentes: verde (agua de lluvia), azul (agua superficial o subterránea) y gris (agua contaminada). Esta métrica permite evaluar la eficiencia hídrica de los cultivos.

**Objetivos:** Este estudio tuvo como objetivo examinar el uso del agua en la agricultura colombiana mediante una revisión sistemática, enfocándose en la gestión del recurso hídrico y su utilidad para orientar decisiones sobre la ubicación óptima de cultivos según sus necesidades de agua.

**Materiales y métodos:** Se realizó una revisión sistemática de 37 estudios científicos sobre huella hídrica agrícola en Colombia, que en conjunto cubren 44 cultivos distribuidos en diferentes regiones del país. Se analizaron los valores reportados de huella hídrica por cultivo y su distribución entre los tres componentes mencionados.

**Resultados:** Los cultivos más estudiados fueron: hortalizas (20%), café y cacao (19%), frutales (17%), flores (12%), cereales (10%) y otros (22%). En promedio, el 72% del agua utilizada corresponde al componente verde, el 15% al azul y el 13% al gris, lo que indica una alta dependencia del agua de lluvia en la agricultura nacional.

**Conclusiones:** Los resultados destacan la necesidad de una planificación hídrica basada en evidencia, que permita mejorar la eficiencia del uso del agua en el sector agrícola colombiano, contribuyendo así a su sostenibilidad.

**Palabras clave:** Huella Hídrica, Agricultura, Agua Virtual, Sostenibilidad, Recurso Hídrico, Demanda hídrica.

### How to cite?

Góngora-Duarte AF, Trujillo-González J M. State of the Art of Agricultural Water Footprint in Colombia. Ingeniería y Competitividad, 2025, 27;(2):e-30214581

<https://doi.org/10.25100/iyc.v27i2.14581>

Recibido: 17-11-24

Evaluado: 4-12-24

Aceptado: 24-02-25

Online: 23-04-25

### Correspondence

andres.gongora@unillanos.edu.co



Spanish version



## Contribution to the literature

### Why was it conducted?


Finding the most important Agricultural Water Footprint studies in the Colombian setting was the primary goal of this research. To accomplish this, a comprehensive analysis was carried out to assess the water needs of agricultural production, taking into account various crops and their respective production methods. The distribution of the Agricultural Water Footprint, the components that were evaluated, and the water footprint in different regions were also compared.

### What were the most relevant results?

Green water footprint, which includes precipitation water that is directly absorbed by crops, accounts for the lion's share of the nation's agricultural water footprint, according to the study's most important findings. The blue water footprint represents irrigation water, and the grey water footprint is the last one. It is worth mentioning that not all documents incorporate all three aspects of the water footprint. For example, some may make reference to chemical fertilizers used in organic agriculture. Since the majority of crops in the country rely on rainwater to meet their water needs, these findings suggest that, generally speaking, there is little competition for water resources. But this may cause ecosystems and other productive activities to compete with humans for water resources.

### What do these results contribute?

These findings shed light on the current status of Colombia's agricultural Water Footprint by comparing the water needs of various agricultural products. Here, data on crop water consumption is compiled for use in agricultural planning, which is an important document for those making decisions about agricultural production systems and water resource management. Lastly, this data inspires more studies in fields where there is a dearth of information about the agricultural water footprint.





## Introduction

Water is a crucial natural resource for societal subsistence and development, serving as a foundation for human progress, the natural environment, and the economy (1,2). Moreover, water resources are essential for agriculture and the quality of life in rural communities (3). Water consumption is estimated to have increased at double the rate of population growth, resulting in various regions experiencing water stress. By 2050, it is projected that 52% of the global population will live in regions experiencing water stress (4). The sustainable management of water resources and the assurance of food security are essential for the advancement of human society (5). The Sustainable Development Goals (SDG) incorporate targets concerning water resources, specifically in SDG 6 “Clean water and sanitation” and SDG 12 “Sustainable production and consumption” (6,7,8). These objectives acknowledge the significance of tackling the escalating issue of water scarcity, alongside the rising demand and strain on water resources (9,10).

Human activities necessitate substantial water resources for development, with agriculture accounting for the majority of usage at 91%, followed by industrial and domestic consumption at 5% and 4%, respectively (11,12). Population and economic growth are driving water resources to unsustainable limits, exacerbated by irrational usage, ecological degradation, and climate change (13,14). At a global scale, the availability of fresh water is problematic due to overexploitation, resulting in environmental issues (15). Recent years have seen an increase in the impact of human activities on natural resources, leading to a focus on the use of environmental indicators to quantify these effects. The aim is to mitigate these impacts and implement comprehensive management of natural resources.

Hoekstra and Hung (16) developed a methodology known as the water footprint to quantify and analyze water consumption, aiming to understand water use in human productive activities and assess the pressure on water resources. The water footprint serves as an indicator assessing the impact of human activities on freshwater resources (16). The water footprint model encompasses both the direct consumption by individuals and the indirect utilization of water resources (17). The water footprint consists of three components: green, blue, and grey. The green water footprint represents the effective precipitation utilized by crops; the blue water footprint pertains to the surface and groundwater resources utilized during production; and the grey water footprint indicates the fresh water required to dilute pollutant loads to achieve natural background concentrations and adhere to environmental regulations (11).

This term denotes the volume of water utilized in the production of a good or service, encompassing the entire supply chain (16). This indicator is essential for advancing the sustainable management of water resources, particularly in the agricultural sector (18). The water footprint serves as a multidimensional indicator reflecting the volume of water consumed by source and the volume affected by contamination, categorized by type. The components of water consumption are delineated in both geographical and temporal contexts (11). This serves as a metric for both consumption volume and water contamination levels. The water footprint does not correlate with the severity of local environmental impacts resulting from water consumption (19,11).



Numerous studies have examined the Water Footprint of agricultural crops to assess the water consumption associated with major crops cultivated in various regions (20,21,22). The concept and its application have been utilized to deliver pertinent information regarding the sustainable utilization of water resources in agricultural production (18). The methodology utilized within agricultural frameworks typically assesses the green and blue water footprint. In certain regions of Latin America, the authors indicate that predominantly green water is utilized for crop development (23), which is the most significant component for agricultural growth. The green water footprint has the largest volume relative to the other types of footprints. In contrast to other factors, there is not always an impetus to minimize it, as it does not create direct competition for water resources. Nonetheless, it suggests competition for territorial use, as evidenced by the expansion of the agricultural frontier (24). The water footprint of crops is typically quantified as yield per unit volume, expressed in cubic meters per ton ( $m^3/ton$ ) or liters per kilogram (L/kg). The water footprint of agricultural crops is defined as the total volume of freshwater utilized, both directly and indirectly, to produce a product. This includes water consumption and pollution associated with each stage of the production chain (11).

The water footprint serves as an indicator of the pressure exerted on water resources, quantified in terms of water volume based on its usage. This necessitates the implementation of instruments that facilitate decision-making for the comprehensive management of water resources, considering the involvement of stakeholders and institutions (25). This review study aims to analyze the concept of Water Footprint as it pertains to agricultural crops, specifically within the Colombian context, with the following objective: Review the state of the art of agricultural water footprint in Colombia, contrasting the water footprint across various crop systems. This approach aims to generate a comprehensive tool for agricultural planning, offering valuable insights into sustainable water use in different production systems.

## Materials and methods

### Data collection

The literature review data were sourced from the Scopus, Google Scholar, and Springer Open databases. An approach was adopted to conduct the review of the water footprint in agricultural crops. The review's scope was delineated, concentrating on the systematic assessment of the water footprint across various crops. The employed keywords were: "Water Footprint" AND "Agriculture" AND "Crops" AND "Water Use". Articles in English and Spanish, published from 2003 to 2024, were selected.

The subsequent phase of the review concentrated on the water footprint of agricultural crops in Colombia. Comprehensive searches were performed in databases including Scopus, Web of Science, and Google Scholar, utilizing keywords such as: "Water Footprint," "Agriculture," "Sustainability," and "Agricultural Crops." References included scientific articles, undergraduate theses, and reports published in the last decade in English and Spanish, concentrating on the application of the water footprint in agricultural crops in Colombia. Using Microsoft Excel, data from every publication was methodically gathered and painstakingly arranged, therefore enabling a thorough and rigorous examination.



## Inclusion and exclusion criteria


The inclusion criteria for this research included studies on the agricultural water footprint in Colombia that used thorough, repeatable methodology. Priority was given to works in Spanish or English that provided extensive and accurate data for comparison and critical analysis. In contrast, research that did not primarily focus on Colombia's agricultural sector, employed faulty or insufficient methodology, or did not provide full access to its data and finding was rejected. Research pertaining to the water footprint of corporate, community, or livestock production was excluded from the review to preserve the emphasis on agricultural crops.

## Results and Discussion

### Agricultural Water Footprint in Colombia

Colombia possesses significant global water resources; however, this resource is jeopardized by inadequate planning, unequal distribution, and climate change, resulting in constraints on water utilization (24,26). In 2020, national water demand comprised 43.25% for agriculture and post-harvest activities, while the remainder was allocated to non-agricultural uses, including hydropower, domestic consumption, livestock, and industry (27). In recent years, numerous studies have been conducted on water in Colombia, primarily concentrating on water resource management, utilization, and quality. These have been conducted in significant regions within sectors characterized by heightened agricultural and industrial activity (24,28).


A total of 37 documents were identified (Fig. 1) that address the calculation of the water footprint of agricultural crops in Colombia. Of these, 47% pertain to undergraduate theses, 45% to scientific articles, and the remaining 8% to institutional reports.



**Figure 1.** Type of publication of Water Footprint of agricultural crops in Colombia.

## Crops


A total of 44 agricultural crops were identified and classified based on their system (Fig. 2). The majority of the publications concentrate on semi-annual crops, comprising 17 documents, with maize, rice and sugar cane as the principal crops. Fifteen perennial crops were documented, with cocoa, coffee, avocado, and cotton being the most prominent. Eight greenhouse crops were identified, including notable varieties such as flowers, tomatoes, and strawberries. Ultimately, one pertinent document was located regarding forest crops.



**Figure 2.** Agricultural cropping systems in water footprint studies in Colombia. Source: Authors.


### Spatial distribution of agricultural crops

A total of 33 study areas were identified across Colombian territory, distributed among various departments. The various crops were categorized collectively. This categorization facilitates enhanced visualization and analysis of the spatial distribution of various crop types (Fig. 3). The Cundinamarca department is distinguished by eight study areas, while Tolima and Boyacá each possess four locations. The departments of Meta, Santander, and Huila each have three study areas, whereas Antioquia has two. Arauca, Nariño, Quindío, Valle del Cauca, and Córdoba each possess a singular study area, indicating the geographic dispersion of Agricultural Water Footprint research in Colombia.



**Figure 3.** Geographical distribution of crops in the reviewed studies. Source: Authors

A total of 37 studies identified 44 crops. Figure.4 illustrates the categorization of crops based on water footprint assessment in Colombia. The category with the greatest number of studies is coffee and cocoa, accounting for 20%, followed by fruit crops at 17%, which includes Hass avocado, tree tomato, and Valencia orange. Vegetables account for 17% of the total, including onion and tomato, whereas flower crops comprise 13%, featuring astromelia flower, carnation, and gerbera. A separate category encompasses certain cereals, specifically rice and maize, while the remaining 23% pertains to crops with fewer studies, including oil palm, tubers, and forest trees.



**Figure 4.** Subdivision of crops analyzed in the reviewed studies. **Source: Authors.**

#### Assessment of the water footprint in agricultural crops

Numerous studies have been conducted in Colombia regarding the Water Footprint associated with agricultural production. Table.1 displays the primary findings of these studies. The investigations align in assessing water consumption in crops to evaluate sustainability concerning the water supply in the study areas (29,30). The study encompasses variables including crop type, climate, precipitation, and soil, primarily focusing on the measurement of the Blue Footprint and Green Footprint in crops, with a secondary emphasis on the grey water footprint. Several authors have estimated the Water Footprint of crops at the basin level (30), whereas others have focused on controlled environments, such as greenhouse crops (31). Conversely, certain crops cultivated through dryland practices exhibit total green water consumption, resulting in a blue water footprint of zero. However, for crops like coffee, the blue water footprint is attributed to processing and post-harvest activities (24).

**Table 1.** Main results of the Agricultural Water Footprint in Colombia

<b>Cultivation system</b>	<b>Crop</b>	<b>Location</b>	<b>GWF</b>	<b>BWF</b>	<b>GWF</b>	<b>Total, WF(m<sup>3</sup>/ton)</b>	<b>Reference</b>	
<b>Perennial crops</b>	Coffee	Villeta	19.996		1.667	21.633	<a href="#">(32)</a>	
	Cotton	Natagaima- Tolima	1.158	749,5	150	2.058	<a href="#">(33)</a>	
		Apartadó (Antioquia)				14.344		
		Tame (Arauca)				15.057		
		Garzón (Huila)				13.475		
		Tumaco (Nariño)				13.719		
		Cocoa	San Vicente (Santander)				22.758	<a href="#">(34)</a>
			Chaparral (Tolima)				23.239	
		Hass Avocado	Quindío	3630	0	1315	4.945	<a href="#">(35)</a>
		Coffee	(Chinchiná, Venecia, El Tambo, Buenavista)	8746		7000	15.749	<a href="#">(36)</a>
		Cocoa	Porce River Basin (Antioquia)	19.428		346	19.346	<a href="#">(37)</a>
		Cocoa	Norte de Santander	13.189	5687		18.876	<a href="#">(38)</a>
		Coffee	Porce River Basin (Antioquia)				7.500	<a href="#">(37)</a>
		Coffee	Huila	6.328	233,2		6.561	<a href="#">(24)</a>
		Dragon fruit	Palestina (Huila)	467	2444,34		2.911	<a href="#">(39)</a>
		Valencia Orange	Puerto López	19735.8	25634.9	77	45488,6	<a href="#">(40)</a>
		Coffee	Anolaima	7.823	9.11	17.93	7.850.5	<a href="#">(41)</a>
	Tree Tomato	Antioquia	248	31	7	286	<a href="#">(37)</a>	
	Oil Palm	Guayuriba River basin, Meta	347,8	141,3		489,1	<a href="#">(42)</a>	
	Tahiti Lime	Angula River basin (Santander)	350	82.88	19.51	452,3	<a href="#">(43)</a>	



Semiannual crops							
Onion	Samacá	65.28	78.7	52.41	196.4		(44)
R12 Potato	El Rosal	169.12	32.80	190.99	392.91		(45)
Spring Onion	Porce River Basin (Antioquia)	402	431	13	846		(37)
Potato	Porce River Basin (Antioquia)	390	78	11	479		(37)
Creole Potato	Porce River Basin (Antioquia)	578	105	16	699		(37)
Onion	Duitama Y Samacá (Boyacá)	151,52	107,3	332,5	591,32		(46)
Potato	Duitama Y Samacá (Boyacá)	200,5	42,93	251,5	494,93		(46)
Rice	Espinal (Tolima)	549,26	184,85	32,02	766,13		(47)
Plantain	Porce River Basin (Antioquia)	2051		39	2.090		(37)
Sugarcane	Villeta	1.856		323	2.179		(32)
Sugarcane	Amaime River Basin (Valle del Cauca)	117	37,44		154.51		(48)
Corn (Maize)	Villeta	2.095		1458.3	3554.2		(32)
Bean	Porce River Basin (Antioquia)	4269		187	4.456		(37)
	Guamo (Tolima)	204,3	1345,6	1033,36	2583,2		
	Cúcuta	179,3	1284,6	1713,24	3177,2		
Rice	La apartada (Córdoba)	362,6	2007,4	1581,55	3951,6		(49)
	Puerto López (Meta)	241,3	1987,3	1900,66	4129,3		
Corn (Maize)	Porce River Basin (Antioquia)	5.696		141	5.837		(37)
Cassava	Villeta	405		277.8	683		(32)
Plantain	Villeta	2.394		547	2.941		(32)
Alstroemeria	Facatativá	1.76	0.95	0.77	3.5		(50)
Flower							
Tomato	Fomeque	87.84	13	233.56	334.4		(51)
Tomato (Greenhouse)	Alto Ricaurte (Boyacá).				1207.4		(52)
Tomato (open field)	Alto Ricaurte (Boyacá).				17411.6		(52)
Gerbera	Facatativá	3.12	1.34	0.82	5.28		(53)
Estatice Flower	Tocancipá	0.62	3.78	1.94	6.34		(54)
Liatris Flower	Tocancipá	1.32	0.91	4.27	6.50		(54)
Strawberry	Sibaté	55.13	22.68	3.84	81.65		(55)
Carnation	Bogotá	2.967	5.476	97.10	8.540		(56)
Forestry	Eucalyptus				1.764		(57)

**GWF:** Green Water Footprint, **HHA:** Blue Water Footprint, **GWF:** Grey Water Footprint **Source:** Authors.

The water footprint values for coffee cultivation are 21,633 m<sup>3</sup>/ton in the Villeta area of Cundinamarca and 6,561 m<sup>3</sup>/ton in the department of Huila. The observed difference is attributable to variations in climatic conditions, including precipitation, and soil types. The first study incorporates the water utilized in the post-harvest process, whereas the second study excludes the grey water footprint (26,17). A study conducted by Bulsink et al. (58) in Indonesia indicates a value of 22,907 m<sup>3</sup>/ton for coffee cultivation, comprising 96% green water and 4% grey water. In contrast, eight study areas have been identified for cocoa cultivation, detailing the water consumption of this crop: Apartadó (Antioquia) at 14,344 m<sup>3</sup>/ton, Tame (Arauca) at 15,057 m<sup>3</sup>/ton, Garzón (Huila) at 13,475 m<sup>3</sup>/ton, Tumaco (Nariño) at 13,719 m<sup>3</sup>/ton, San Vicente (Santander) at 22,758 m<sup>3</sup>/ton, and Chaparral (Tolima) at 23,239 m<sup>3</sup>/ton (34). Additional research indicates values in the Porcè River basin, Antioquia, and Norte de Santander, recorded at 19,346 m<sup>3</sup>/ton and 18,876 m<sup>3</sup>/ton, respectively (37,38).

The water footprint of cocoa cultivation in Ecuador is estimated at 24,602.70 m<sup>3</sup>/ton. The disparity arises from the utilization of 30.22% blue water and 69.78% green water for crop development in Ecuador (23). In Colombia, water consumption for the crop is derived solely from green water, with no reported values for grey water. In the context of cotton cultivation, the Natagaima area in Tolima reports a value of 2,058 m<sup>3</sup>/ton, indicating a substantial blue water consumption of 36.41%. This underscores the necessity of irrigation water for crop development. Chapagain and Hoekstra (59) conducted a study assessing the total water footprint associated with cotton cultivation across multiple countries. The results revealed values of 1,534 m<sup>3</sup>/ton in Greece, 1,325 m<sup>3</sup>/ton in Spain, and 2,320 m<sup>3</sup>/ton in Turkey. The reported value for Colombia is higher; however, it does not indicate a significant difference when compared to the regions analyzed. The observed difference in values may result from the implementation of more efficient irrigation systems, as indicated by the total value of the blue water footprint (60).


A study on Hass Avocado cultivation indicates a water usage of 4,945 m<sup>3</sup>/ton, comprising 73.4% green water and 26.6% grey water associated with Nitrogen contamination. Importantly, in Colombia, this crop does not require supplemental irrigation (35). Comparable research, including that conducted by Cerna and Olivares (61) in Peru, indicates a value of 956 m<sup>3</sup>/ton. Conversely, in Chile, the average water footprint of Avocado is 389 m<sup>3</sup>/ton (62). A notable difference is evident in the value obtained in Colombia, suggesting variations in the estimation of the water footprint related to the crop's development time. Sugarcane cultivation encompasses two study areas, revealing reported values of 2,197 m<sup>3</sup>/ton for Villeta Cundinamarca and 154.51 m<sup>3</sup>/ton in the Amaime River basin in Valle del Cauca. The water requirements of the two areas differ due to variations in climatic conditions, soil types, and the timeframes utilized in calculating the water footprint (32,48). In Brazil, sugarcane cultivation yields values of 98 m<sup>3</sup>/ton in dry land conditions and 119 m<sup>3</sup>/ton under irrigation (63).

Tomato cultivation occurs in two modalities: open field and greenhouse. The water consumption values are 334.4 m<sup>3</sup>/ton for open field and 1,207 m<sup>3</sup>/ton for greenhouse cultivation. This indicates a significant difference in green and blue water usage, with greenhouse crops requiring a higher volume of blue water throughout their production cycle (52). Mekonnen and Hoekstra (64) report an average water footprint of 214 m<sup>3</sup>/ton for tomatoes. Furthermore, in regions like Northwest São

Paulo, Brazil, the average water footprint for tomatoes is reported to be 94 m<sup>3</sup>/ton, comprising 72% green water consumption, 20% blue water, and 8% grey water (65).

### Colombian agricultural water footprint colors

Figure 5 illustrates the components examined in the analyzed studies. Sixty percent of these encompass the three components (green, blue, and grey), while twenty-five percent focus solely on the green and grey footprints. The primary contributor to the water footprint in agricultural crops is green water, which is influenced by seasonality and precipitation (5). Most studies on grey water footprint estimates have focused on nitrogen (N) and phosphorus (P) pollution (66). It is noted that 12% of the studies focus exclusively on green and blue water.




**Figure 5.** Subdivision according to the components considered in the studies **Source:** Authors.

### Distribution of the agricultural water footprint

Studies indicate that green water accounts for the highest water consumption at 72%, followed by blue water at 15%, and grey water at 13% (Fig.6). The National Water Study 2022 indicates that the green water footprint has the highest participation. This is due to the significant rainfall levels, which decrease reliance on irrigation water and lower the opportunity cost of water use in Colombian agriculture (27). The blue water footprint is relatively low, as not all agricultural crops are cultivated using irrigation practices. This value reflects crops with significant water requirements that exceed the available water resources in regions such as Tolima, Huila, and Magdalena (67).

The grey water footprint represents the component with the least contribution. The findings suggest that not all documents disclose their value as a result of employing organic farming practices (42). The calculation of this footprint necessitates specific data, including soil texture, yield, leaching, and environmental factors. This can also be mitigated through precision farming practices and soil conservation (68).



**Figure 6.** Distribution of the agricultural water footprint in Colombia. **Source:** Authors.

The sustainable management of water resources is a primary focus of Colombia's public policies, especially in sectors like agriculture. The 2018 CONPES 3934 document, entitled "Green Growth Policy," emphasizes the necessity of implementing production practices that safeguard this essential resource, meeting present demands without jeopardizing its future availability (69). Nonetheless, inconsistencies between environmental and agricultural policies endure, resulting in substantial deficiencies in the shift towards sustainable production models, as observed by Forero and González (70). Current policies predominantly emphasize the oversight of water usage; however, a more profound integration of water supply and demand in critical productive sectors is necessary. Moreover, instruments such as the National Climate Change Adaptation Plan (PNACC) and Law 1931 of 2018 advocate for water adaptation measures but encounter obstacles including inadequate institutional coordination, constrained funding, and limited local engagement (71). This highlights the necessity to enhance territorial planning and improve water governance that effectively integrates public policies with rural development and environmental conservation.

Since 1998, the National Water Studies (NWS) (Estudios Nacionales del Agua ENA) conducted by IDEAM have emerged as a crucial instrument for water resource management in Colombia. The 2014 national study introduced the concept of water footprint, facilitating an analysis of water consumption in agriculture. Key crops such as coffee, sugarcane, and oil palm were emphasized, indicating that agriculture in Colombia predominantly relies on green water, with minimal dependence on blue water or irrigation. The ratio of green to blue water footprints differs between regions and hydrographic subzones (72).

The 2018 NWS estimated the water demand and blue and green water footprints of Colombia's agricultural sector by quantifying crop water requirements and assessing the available water sources in each region. This resulted in the development of national maps depicting green and blue water footprints, categorized by hydrographic subzones. The analyzes highlighted the significant water consumption of crops including irrigated rice, plantain, and oil palm (73).

In its 2022 NWS, IDEAM subsequently introduced the Basin Water Return Index (IARC), which pertains to the water footprint. This indicator demonstrates the ability of hydrographic subzones to support water usage stemming from activities such as agriculture. Furthermore, the Water Pressure Index on Ecosystems (IPHE) was integrated. This index evaluates the green water footprint of the agricultural and forestry sectors within a hydrographic subzone, juxtaposing it with the availability of green water during the corresponding timeframe. The sustainability of the agricultural sector at the national level was evaluated through this index. The research additionally encompassed forecasts of water usage by productive sectors until 2040 (27).

The National Water Studies have served as essential instruments for national public policy, employing the Water Footprint as a criterion to assess agricultural water consumption. These studies examined water usage linked to different cropping systems, producing comprehensive maps that illustrate each element of the Water Footprint. They presented a lucid depiction of the strain agricultural production exerted on water resources, thereby establishing a robust basis for sustainable water management within the agricultural sector (27).

## Conclusions

The review permitted the establishment of the current state of the art in terms of agricultural water footprint in Colombia. The majority of bibliographic production is for undergraduate and postgraduate theses, which may indicate a need to review the precision of the results and methodologies used, particularly when compared to scientific research in the same field.

It was discovered that the water footprint of various crops has been quantified in different regions, with the green water footprint accounting for the majority of the total water footprint. This is due to the fact that the main crops are grown using rainfed practices in areas with high rainfall rates. Agriculture stands out as an activity that captures and uses rainwater, which accounts for a significant portion of agricultural production in Colombia. Although the green water footprint does not imply direct competition for water consumption, it does result in competition for land use as the national agricultural frontier expands.

In Colombia, using rainwater in agriculture reduces the need for irrigation while also promoting sustainability. It is recommended to use efficient technologies that prioritize this source, in line with SDG 12. Furthermore, assessing the water footprint contributes to reducing fertilizer contamination and preserving water quality, which aligns with SDG 6. Understanding water requirements and availability is critical for avoiding resource constraints and promoting sustainable use, emphasizing the need for future research in agricultural water management.

The agricultural sector in Colombia necessitates the implementation of policies that emphasize the sustainable utilization of water, thereby ensuring a harmonious equilibrium between production requirements and the conservation of environmental resources.



## CrediT authorship contribution statement

Conceptualization - Ideas: Andrés Felipe Góngora Duarte, Juan Manuel Trujillo Gonzalez. Data Curation: Andrés Felipe Góngora Duarte. Formal analysis: Andrés Felipe Góngora Duarte. Investigation: Andrés Felipe Góngora Duarte. Methodology: Andrés Felipe Góngora Duarte. Project Management: Andrés Felipe Góngora Duarte. Resources: Andrés Felipe Góngora Duarte, Supervision: Juan Manuel Trujillo González. Validation: Andrés Felipe Góngora Duarte, Juan Manuel Trujillo González. Writing - original draft - Preparation: Andrés Felipe Góngora Duarte, Juan Manuel Trujillo González. Writing - revision and editing - Preparation: Juan Manuel Trujillo González.

Financing: does not declare. Conflict of interest: does not declare. Ethical aspect: does not declare.

## References

1. Jie H, Khan I, Alharthi M, Zafar MW, Saeed A. Sustainable energy policy, socio-economic development, and ecological footprint: The economic significance of natural resources, population growth, and industrial development. *Utilities Policy*. 2023;81. Available in: <https://doi.org/10.1016/j.jup.2023.101490>
2. Abu-Zeid MA, Abu-Zeid KM, Halim MK. Impact of Climate Change on Water Resources. In: *Climate Changes Impacts on Aquatic Environment: Assessment, Adaptation, Mitigation, and Road Map for Sustainable Development*. Cham: Springer Nature Switzerland; 2025. p. 121-157. Available in: [https://doi.org/10.1007/978-3-031-74897-4\\_5](https://doi.org/10.1007/978-3-031-74897-4_5)
3. Ardila C, del Pilar M. Gestión del agua para riego en prácticas de agricultura desde la complejidad ambiental [specialization thesis]. Bogotá: Fundación Universidad de América; 2018. Available in: <https://repository.uamerica.edu.co/server/api/core/bitstreams/2ca23716-2b39-4599-8764-b2781e1f364b/content>
4. Organización de las Naciones Unidas para la Educación, la Ciencia y la Cultura. Agua y cambio climático [Internet]. UNESCO; 2020 [accessed: December 10, 2024]. Available in: <https://unesdoc.unesco.org/ark:/48223/pf0000373611>
5. Makanda K, Nzama S, Kanyerere T. Assessing the role of water resources protection practice for sustainable water resources management: a review. *Water*. 2022;14(19):3153. Available in: <https://doi.org/10.3390/w14193153>
6. United Nations. Sustainable Development Goal 6 Synthesis Report 2018 on Water and Sanitation [Internet]. New York: United Nations; 2018 [accessed: December 10, 2024]. Available from: [https://sustainabledevelopment.un.org/content/documents/19901SDG6\\_SR2018\\_web\\_3.pdf](https://sustainabledevelopment.un.org/content/documents/19901SDG6_SR2018_web_3.pdf)
7. Verma N, Talwar P, Upadhyay A, Singh R, Lindenberger C, Pareek N, et al. Food-Energy-Water Nexus in compliance with Sustainable Development Goals for integrating and managing the core environmental verticals for sustainable energy and circular economy. *Sci Total Environ*. 2024;930:172649. Available in: <https://doi.org/10.1016/j.scitotenv.2024.172649>
8. Papamichael I, Voukkali I, Stylianou M, et al. Sustainable production and consumption. *Euro-Mediterr J Environ Integr*. 2024;9:2003–2008. Available in: <https://doi.org/10.1007/s41207-024-00594-0>
9. Miao J, Song X, Zhong F, Huang C. Sustainable development goal 6 assessment and attribution analysis of underdeveloped small regions using integrated multisource data. *Remote Sens*. 2023;15(15):3885. Available in: <https://doi.org/10.3390/rs15153885>
10. Ingrao C, Strippoli R, Lagioia G, Huisingh D. Water scarcity in agriculture: An overview of causes, impacts and approaches for reducing the risks. *Heliyon*. 2023;9(8). Available in: <https://doi.org/10.1016/j.heliyon.2023.e18507>



11. Hoekstra AY, Chapagain AK, Aldaya MM, Mekonnen MM. The Water Footprint Assessment Manual. February 2011. Available in: <http://doi.org/978-1-84971-279-8>
12. Lovarelli D, Bacenetti J, Fiala M. Water footprint of crop productions: A review. *Science of the Total Environment*. 2016;548:236-251. Available in: <https://doi.org/10.1016/j.scitotenv.2016.01.022>
13. Hua E, Han X, Engel BA, Guan J, Sun S, Wu P, et al. Developing a sustainable assessment framework for identifying industrial water suitability Perspective on the water-energy-food nexus. *Agricultural Systems*. 2024;220:104065. Available in: <https://doi.org/10.1016/j.agsy.2024.104065>
14. Peng Q, He W, Kong Y, Shen J, Yuan L, Ramsey TS. Spatio-temporal analysis of water sustainability of cities in the Yangtze River Economic Belt based on the perspectives of quantity-quality-benefit. *Ecological Indicators*. 2024;160:111909. Available in: <https://doi.org/10.1016/j.ecolind.2024.111909>
15. Zhang GP, Hoekstra AY, Mathews RE. Water Footprint Assessment (WFA) for better water governance and sustainable development. *Water resources and industry*. 2013;1-2:1-6. Available from: <http://dx.doi.org/10.1016/j.wri.2013.06.004>
16. Hoekstra AY, Hung PQ. Virtual water trade: a quantification of virtual water flows between nations in relation to international crop trade. Value of Water Research Report Series No. 11. UNESCO-IHE; 2002. Available in: <https://www.waterfootprint.org/resources/Report11.pdf>
17. Hoekstra AY, Wiedmann TO. Humanity's unsustainable environmental footprint. *Science*. 2014;344(6188):1114-1117. Available from: <https://doi.org/10.1126/science.1248365>
18. Dehghanpir S, Bazrafshan O, Ramezani Etedali H, Holisaz A, Collins B. Water scarcity assessment in Iran's agricultural sector using the water footprint concept. *Environment, Development and Sustainability*. 2023;1-26. Available in: <https://doi.org/10.1007/s10668-023-03852-3>
19. Hoekstra AY, Hung PQ. Globalization of water resources: International virtual water flows in relation to crop trade. *Global Environmental Change*. 2005;15(1):45-56. Available in: <https://doi.org/10.1016/j.gloenvcha.2004.06.004>
20. Finogenova N, Dolganova I, Berger M, Núñez M, Blizniukova D, Müller-Frank A, et al. Water footprint of German agricultural imports: local impacts due to global trade flows in a fifteen-year perspective. *Science of the Total Environment*. 2019;662:521-529. Available in: <https://doi.org/10.1016/j.scitotenv.2019.01.264>
21. Liu L, Hu X, Zhan Y, Sun Z, Zhang Q. China's dietary changes would increase agricultural blue and green water footprint. *Science of the Total Environment*. 2023;903:165763. Available from: <https://doi.org/10.1016/j.scitotenv.2023.165763>
22. Garrido-Rubio J, González-Piqueras J, Calera A, Osann A. Remote sensing-based green and blue agricultural water footprint estimation at the river basin scale. *Ecological Indicators*. 2024;167:112643. Available in: <https://doi.org/10.1016/j.ecolind.2024.112643>
23. Dumes JMM, Peralta SLP. Valor económico de la huella hídrica de cacao arriba (Theobroma Cacao). *Revista de la Universidad del Zulia*. 2023;14(41):94-130. Available in: <https://doi.org/10.46925//rdluz.41.06>
24. Camacho WA, Uribe DA. Estimación de la huella hídrica azul y verde de la producción cafetera en ocho cuencas en el sur del departamento del Huila. *Revista de investigación agraria y ambiental*. 2018;9(2):338-354. Available in: <https://doi.org/10.22490/21456453.2284>
25. Al-Saidi M. Conflicts and security in integrated water resources management. *Environmental Science & Policy*. 2017;73:38-44. Available in: <https://doi.org/10.1016/j.envsci.2017.03.015>
26. Ordóñez JI. El agua y el sector rural en Colombia. *Revista de ingeniería*. 2020;(49):10-17. Available in: <https://revistas.uniandes.edu.co/index.php/rdi/article/download/7486/7883>
27. Instituto de Hidrología, Meteorología y Estudios Ambientales (IDEAM). Estudio nacional del agua 2022. IDEAM; 2022. Available in: [https://www.andi.com.co/Uploads/ENA%202022\\_compressed.pdf](https://www.andi.com.co/Uploads/ENA%202022_compressed.pdf)
28. Rincón MAP, Hurtado IC, Restrepo S, Bonilla SP, Calderón H, Ramírez A. Water footprint measure method for tilapia, cachama and trout production: study cases to Valle del Cauca (Colombia). *Ingeniería y competitividad*. 2017;19(2):109-120. Available in: <https://doi.org/10.25100/iyc.v19i2.5298>



29. Tovar-Hernández, N. A., Trujillo-González, J. M., Muñoz-Yáñez, S. I., Torres-Mora, M. A., & Zárate, E. (2017). Evaluación de la sostenibilidad de los cultivos de arroz y palma de aceite en la cuenca del río Guayuriba (Meta, Colombia), a través de la evaluación de huella hídrica. *Orinoquia*, 21(1), 52-63. <https://doi.org/10.22579/20112629.394>
30. Vargas-Pineda OI, Trujillo-González JM, Torres-Mora MA. Supply-demand of water resource of a basin with high anthropic pressure: case study Quenane-Quenanito Basin in Colombia. *Air, Soil and Water Research*. 2020;13:1178622120917725. Available in: <https://doi.org/10.1177/1178622120917725>
31. Bojacá CR, Wyckhuys KA, Schrevels E. Life cycle assessment of Colombian greenhouse tomato production based on farmer-level survey data. *Journal of Cleaner Production*. 2014;69:26-33. Available from: <https://doi.org/10.1016/j.jclepro.2014.01.078>
32. Ortiz Caballero LD. Cálculo de la huella hídrica de los sistemas productivos agrícolas de la microcuenca Quebrada Cune, municipio de Villeta, Cundinamarca [Undergraduate thesis]. Bogotá: Universidad de La Salle; 2014. Available in: <https://ciencia.lasalle.edu.co/bitstreams/6cf6f33b-397a-42a6-a193-810184f56bbc/download>
33. Díaz Fonseca OD, Lizarazo Ariza JA. Determinar la huella hídrica del cultivo de algodón (*Gossypium*) de la empresa Coagronat Ltda. en el municipio Natagaima-Tolima [Undergraduate thesis]. Bogotá: Universidad de La Salle; 2016. Available in: <https://ciencia.lasalle.edu.co/bitstreams/dd48c0c9-ea94-4e6c-8cf1-646250de3014/download>
34. Ortiz-Rodríguez OO, Naranjo CA, García-Caceres RG, Villamizar-Gallardo RA. Water footprint assessment of the Colombian cocoa production. *Revista Brasileira de Engenharia Agrícola e Ambiental*. 2015;19(9):823-828. Available in: <http://dx.doi.org/10.1590/1807-1929/agriambi.v19n9p823-828>
35. Naranjo JF, Reyes H. Huella hídrica del cultivo de aguacate cv. Hass (*Persea americana* Mill.), en el Distrito de Conservación de Suelos Barbas-Bremen, Quindío, Colombia. *Entre Ciencia e Ingeniería*. 2021;15(29):63-70. Available from: <https://doi.org/10.31908/19098367.1813>
36. Leal-Echeverri JC, Tobón C. The water footprint of coffee production in Colombia. *Revista Facultad Nacional de Agronomía Medellín*. 2021;74(3):9685-9697. Available in: <https://doi.org/10.15446/rfnam.v74n3.91461>
37. Builes Cédula ED. Cuantificación y análisis de sostenibilidad ambiental de la huella hídrica agrícola y pecuaria de la cuenca del río Porce [master's thesis]. Medellín: Universidad Nacional de Colombia; 2013. Available in: <https://repositorio.unal.edu.co/handle/unal/20267>
38. Naranjo-Merino CA, Ortiz-Rodríguez OO, Villamizar-G RA. Assessing green and blue water footprints in the supply chain of cocoa production: A case study in the Northeast of Colombia. *Sustainability*. 2017;10(1):38. Available in: <https://doi.org/10.3390/su10010038>
39. Bautista EAS, Roa JRV, Ortega JAT. Estimación de la huella hídrica para un cultivo de pitahaya amarilla (*Selenicereus megalanthus*). *Publicaciones e investigación*. 2015;9:135-146. Available from: <https://doi.org/10.22490/25394088.1439>
40. Sabogal Ángel JS. Evaluación de la huella hídrica para el cultivo de naranja (Valencia) en un predio del municipio de Puerto López–Meta [master's thesis]. Bogotá: Universidad de La Salle; 2015. Available in: <https://ciencia.lasalle.edu.co/server/api/core/bitstreams/9d02d8f9-6f71-4f96-9b24-5dd14826ba9d/content>
41. Forero Peña AM, Pastor Silva EP. Aproximación a la huella hídrica del cultivo de café en el municipio de Anolaima y determinación de la presión generada por la demanda hídrica y descarga de aguas mieles [undergraduate thesis]. Bogotá: Universidad de La Salle; 2014. Available in: <https://ciencia.lasalle.edu.co/server/api/core/bitstreams/2e3431bc-f008-4322-8601-33cfc6387c4b/content>
42. Trujillo-González JM, Tovar-Hernández NA, Torres-Mora MA. Water footprint of fresh fruit bunches (FFBs) of oil palm in the Piedemonte Llanero, Colombia. *inycomp* [. 2022 Jan. 15 [cited 2024 Nov. 17];24(1). Available in: <https://doi.org/10.25100/iyv.v24i1.10947>
43. Arenas-Jiménez CF, Correa-Torres SN, Pineda-Vargas SM. Estimación de la huella hídrica en la producción agrícola de lima Tahití en la Cuenca La Angula, Santander, Colombia. *Investig Cienc*. 2020;28(79):52-61. Available in: <https://doi.org/10.33064/iyvuaa2020792939>



44. Cely-Reyes GE, Camargo-Guerrero FDR, Fernández-Pérez CJ, Walteros-Torres IY, Serrano-Cely PA. Determination of the water footprint in the production of the bulb onion crop under two irrigation systems in Samaca (Colombia). *Rev Colomb Cienc Hort*. 2023. Available in: <https://doi.org/10.17584/rcch.2023v17i2.16010>
45. Amézquita Bonilla VV, Motta López DF. Evaluación de huella hídrica en la producción de un cultivo de papa R12 ubicado en el municipio El Rosal, Cundinamarca [Undergraduate thesis]. Bogotá: Universidad de La Salle; 2015. Available in: <https://ciencia.lasalle.edu.co/server/api/core/bitstreams/99b5dfb4-eb2b-41f8-9406-755b3a37bc6b/content>
46. Higuera C, Jaimes H. Evaluación de indicadores de huella hídrica en la producción de un cultivo de cebolla de bulbo y papa en los municipios de Duitama y Samacá, Boyacá [specialization thesis]. Bogotá: Universidad Católica de Colombia; 2019. Available in: <https://repository.ucatolica.edu.co/server/api/core/bitstreams/83f56d69-2835-4477-9c8a-8c20d0aed53b/content>
47. Lozano Chamorro D, Cortés González N. Evaluación de la huella hídrica del proceso productivo del arroz (*Oryza sativa*) en el municipio del Espinal–Tolima y su incidencia ambiental en el área de influencia [Undergraduate thesis]. Bogotá: Universidad de La Salle; 2016. Available in: <https://ciencia.lasalle.edu.co/server/api/core/bitstreams/b9b0259c-bf2a-49b0-92a6-7b08fc5134c6/content>
48. Ramírez L, Becerra D, Mora C. Huella hídrica verde y azul de la producción de caña de azúcar orgánica en la zona centro del Valle del Cauca. *Inycomp*. 2021;24(2-2022). Available in: <https://doi.org/10.25100/iyc.v24i02.11264>
49. Murcia Rojas AV, Casadiego Hernández D. Cálculo comparativo de la huella hídrica del sistema productivo de arroz de riego en cuatro zonas arroceras de Colombia por medio del software Cropwat 8.0 [Undergraduate thesis]. Bogotá: Universidad de La Salle; 2018. Available in: <https://ciencia.lasalle.edu.co/server/api/core/bitstreams/42dd5be7-5b05-47db-851a-b29de56c7a08/content>
50. Velandia Rubiano IL, López Romero KJ. Evaluación de la huella hídrica para la producción de flores astromelias (*Alstroemeria* sp) en un predio del municipio de Facatativá, Cundinamarca [undergraduate thesis]. Bogotá: Universidad de La Salle; 2014. Available in: <https://ciencia.lasalle.edu.co/server/api/core/bitstreams/8ce28568-d3bb-48b1-9169-a46cb3d117dc/content>
51. Cruz González M. Análisis y evaluación de la huella hídrica de un cultivo de tomate (*Lycopersicon esculentum*) en el municipio de Fómeque, departamento de Cundinamarca [undergraduate thesis]. Bogotá: Universidad Militar Nueva Granada; 2017. Available in: <https://repository.umng.edu.co/server/api/core/bitstreams/a87279b7-7e2e-4d3c-9fd4-01e29c253aee/content>
52. Prias Gómez AJ. Análisis de huella hídrica de los sistemas productivos de tomate a campo abierto y bajo invernadero en Colombia [master's thesis]. Bogotá: Universidad Jorge Tadeo Lozano; 2015. Available in: <https://expeditiorepositorio.utadeo.edu.co/bitstream/handle/20.500.12010/1767/T065%20%281%29.pdf?sequence=1>
53. Rodríguez Martínez LM, Juya Roa EJ. Determinación y evaluación de la huella hídrica en la producción de un cultivo de gerbera ubicado en el municipio de Facatativá [Undergraduate thesis]. Bogotá: Universidad de La Salle; 2015. Available in: <https://ciencia.lasalle.edu.co/server/api/core/bitstreams/45a94bdc-cba2-40be-b129-91b12a6cd283/content>
54. Rincón Loaiza AM. Determinación de la huella hídrica de flores *Liatris* y *Estatice* en un floricultivo en el municipio de Tocancipá [Undergraduate thesis]. Bogotá: Universidad de La Salle; 2014. Available in: <https://ciencia.lasalle.edu.co/server/api/core/bitstreams/07701673-3825-4674-8b42-4ea68d1e6f22/content>
55. Linares Castillo LM, Melo Rojas L. Evaluación de la huella hídrica del cultivo de fresa en una finca del municipio de Sibaté, Cundinamarca [Undergraduate thesis]. Bogotá: Universidad de La Salle; 2015. Available in: <https://ciencia.lasalle.edu.co/bitstreams/ee2258b0-fbc7-4de4-8362-935162f18903/download>



56. Baez Silva WR. Evaluación de la huella hídrica en la producción de clavel estándar (*Dianthus caryophyllus*) en la sabana de Bogotá [Undergraduate thesis]. Bogotá: Universidad Nacional de Colombia; 2013 [citado el 2025 Mar 11]. Available in: <https://core.ac.uk/download/pdf/198441829.pdf>
57. Vásquez Velásquez G. Los bosques plantados y la huella hídrica, la respuesta hidrológica y la hidrosolidaridad. Departamento de Ciencias Forestales; 2011. Available in: [https://repositorio.unal.edu.co/bitstream/handle/unal/57650/Pantaciones%20y%20Bosques\\_Doc\\_Sectorial\\_Forestal.pdf?sequence=1](https://repositorio.unal.edu.co/bitstream/handle/unal/57650/Pantaciones%20y%20Bosques_Doc_Sectorial_Forestal.pdf?sequence=1)
58. Bulsink F, Hoekstra AY, Booij MJ. The water footprint of Indonesian provinces related to the consumption of crop products. *Hydrol Earth Syst Sci Discuss.* 2010;6(4):5115–37. Available in: <https://doi.org/10.5194/hess-14-119-2010>
59. Chapagain AK, Hoekstra AY, Savenije HHG, Gautam R. The water footprint of cotton consumption: an assessment of the impact of worldwide consumption of cotton products on the water resources in the cotton producing countries. *Ecol Econ.* 2006;60:186–203. Available in: <http://dx.doi.org/10.1016/j.ecolecon.2005.11.027>
60. Tsakmakis ID, Zoidou M, Gikas GD, Sylaios GK. Impact of irrigation technologies and strategies on cotton water footprint using AquaCrop and CROPWAT models. *Environ Process.* 2018;5:181–199. Available in: <https://doi.org/10.1007/s40710-018-0289-4>
61. Cerna Angeles MF, Olivares Hipolito IR. Huella hídrica del arándano azul (*Vaccinium corymbosum*), espárrago verde (*Asparagus officinalis*), palta Hass (*Persea americana*) cultivados en La Libertad, Perú [Undergraduate thesis]. Trujillo: Universidad Nacional de Trujillo; 2018. Available in: <https://dspace.unitru.edu.pe/server/api/core/bitstreams/36a74446-ed45-4c66-884f-db4b184ce33d/content>
62. Panez-Pinto A, Mansilla-Quiñones P, Moreira-Muñoz A. Agua, tierra y fractura sociometabólica del agronegocio. *Actividad frutícola en Petorca, Chile. Bitácora Urbano Territorial.* 2018;28(3):153-60. Available in: <https://doi.org/10.15446/bitacora.v28n3.72210>
63. Scarpore FV, Hernandez TAD, Ruiz-Corrêa ST, Kolln OT, Gava GJ de C, dos Santos LNS, et al. Sugarcane water footprint under different management practices in Brazil: Tietê/Jacaré watershed assessment. *J Clean Prod.* 2016;112(Part 5):4576-84. Available in: <https://doi.org/10.1016/j.jclepro.2015.05.107>
64. Mekonnen MM, Hoekstra AY. The green, blue and grey water footprint of crops and derived crop products. Twente Water Centre, University of Twente, Enschede, The Netherlands. 2011. Available in: <https://doi.org/10.5194/hess-15-1577-2011>
65. Martins Filho JB, Cremasco CP, Noris FJ, dos Santos RM, dos Santos JSP, Dal Pai A, et al. Demanda hídrica do tomate de mesa no município de Paranapuã-SP em diferentes épocas de plantio. *Research, Society and Development.* 2022;11(6). Available in: <https://doi.org/10.33448/rsd-v11i6.28797>
66. Aldaya MM, Rodriguez CI, Fernandez-Poulussen A, Merchan D, Beriain MJ, Llamas R. Grey water footprint as an indicator for diffuse nitrogen pollution: The case of Navarra, Spain. *Science of the Total Environment.* 2020;698:134338. Available in: <https://doi.org/10.1016/j.scitotenv.2019.134338>
67. Arévalo D, Lozano J, Sabogal J. Estudio nacional de huella hídrica Colombia sector agrícola. *Revista Internacional de Sostenibilidad, Tecnología y Humanismo.* 2011;(6):101-126. Available in: <https://upcommons.upc.edu/handle/2099/11915>
68. Borsato E, Galindo A, Tarolli P, Sartori L, Marinello F. Evaluation of the grey water footprint comparing the indirect effects of different agricultural practices. *Sustainability.* 2018;10(11):3992. Available in: <https://doi.org/10.3390/su10113992>
69. Departamento Nacional de Planeación (DNP). CONPES 3934: Política de Crecimiento Verde. Bogotá: Departamento Nacional de Planeación; 2018. Available in: <https://colaboracion.dnp.gov.co/cdt/conpes/econ%3%B3micos/3934.pdf> (Accessed December 5, 2024)
70. Forero N, González C. Agricultura Climáticamente Inteligente (ACI) en Colombia: diagnóstico y retos de política pública. 2020. Available in: <https://www.repository.fedesarrollo.org>



- [co/bitstream/handle/11445/4053/Co Eco Diciembre 2020 Forero y Gonz%C3%A1lez.pdf?sequence=4&isAllowed=y](https://co/bitstream/handle/11445/4053/Co_Eco_Diciembre_2020_Forero_y_Gonz%C3%A1lez.pdf?sequence=4&isAllowed=y) (Accessed December 5, 2024)
71. MinAmbiente National Adaptation Plan (2011). Available in [https://www.minambiente.gov.co/images/cambioclimatico/pdf/Plan\\_nacional\\_de\\_adaptacion/Proyecto\\_nacional\\_de\\_adaptacion\\_al\\_cambio\\_climatico\\_INAP.pdf](https://www.minambiente.gov.co/images/cambioclimatico/pdf/Plan_nacional_de_adaptacion/Proyecto_nacional_de_adaptacion_al_cambio_climatico_INAP.pdf) (accessed December 6, 2024).
  72. IDEAM. Estudio Nacional del Agua 2014. Bogotá D.C.: Instituto de Hidrología, Meteorología y Estudios Ambientales (IDEAM); 2014. Available in: [https://www.andi.com.co/Uploads/ENA\\_2014.pdf](https://www.andi.com.co/Uploads/ENA_2014.pdf) (accessed: December 10, 2024)
  73. IDEAM. Estudio Nacional del Agua 2018. Bogotá D.C.: Instituto de Hidrología, Meteorología y Estudios Ambientales (IDEAM); 2018. Available in: [https://www.andi.com.co/uploads/ena\\_2018-comprimido.pdf](https://www.andi.com.co/uploads/ena_2018-comprimido.pdf) (accessed: December 10, 2024)